



11 Park Drive, Suite 12
Boston, MA 02215

Mouse Anti-Human VEGF-A

ORDERING INFORMATION

Catalog Number:	101-M26
Size:	100 µg
Formulation:	Monoclonal Antibody; Lyophilized
Clone/AB feature:	(#EIC)
Antigen:	Recombinant human soluble Flt-1 (D5)
Application:	ELISA, WB, IP
Stabilizer:	None
Buffer:	PBS pH 7.4 w/o preservative

Description:

Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVE supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA.

Reconstitution:

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Stability:

The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 °C. Frozen aliquots are stable for at least 6 months when stored at -20 °C. **Avoid repeated freeze-thaw cycles!**

Optimal dilutions should be determined by each laboratory for each application.

The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users!

This product is sold for Research Use Only !

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com