

hTert-GFP Mouse Glomerular Podocytes

ORDER INFORMATION

Name of Cells: hTert-GFP Mouse Glomerular Podocytes (hTert-GFP-mGloPods)
Catalogue Number: **cAP-m0024-hTert-GFP**
Product Format: Frozen Vial
Cell Number: > 5 x 10⁵/vial

General Information


Mouse Glomerular Podocytes (mGloPods) were isolated from kidneys of 4-week old C57/Black6 mice and hTert-GFP-mGloPods were selected from mGloPods infected with hTert and GFP expressing lentiviruses with puromycin. A clone of hTert-GFP-mGloPods which maintain the foot process formation were isolated and supplied in frozen vials. Podocyte Growth Medium (cAP-44) is recommended for the expansion of hTert-GFP-mGloPods and the maintaining of foot process formation in vitro. And we warranty a minimum of 30 passages of the cells if cultured following the detailed protocol described below.

Characterization:

Positive: Nephrin, CD2AP abd

Negative: CD31

hTert-GFP-mGloPods are tested negative for common animal pathogen and mycoplasma.



Transformed Mouse Glomerular Podocytes Maintain their Foot Processes in Vitro

Only.

Product Use: hTert-GFP-mGloPods are for Research Use

Shipping: Frozen Vial with > 5 x 10⁵ cells/vial

Handling of Arriving Cells

When you receive the cells in a frozen vial, you can transfer the vial of cells into a -80°C freezer for short term storage or a liquid nitrogen tank for long term storage. Thaw the cells in a 37°C water bath, and then transfer the cells into a T25 flask Quick Coating Solution (cAP-01) pre-coated flasks as described in details in Subculture Protocol.

Subculture Protocol

Pre-coating of T25 flasks: Add 2ml of the Quick Coating Solution Coating Solution (cAP-01) into one T25 flask and make sure the whole surface of the flask is covered with the Coating Solution; Leave the T25 flask with the Quick Coating Solution for 5 minutes; Aspirate off the Coating Solution after 5 minutes and the flask is ready to be used.

- When Cells are nearly confluent, rinse the cells in T25 flask with 5ml HBSS (**Room Temperature, RT**) twice.
- Add 2ml of Trypsin/EDTA (**RT**) (cAP-23) into one T25 flask (make sure the whole surface of the T25 flask is covered with Trypsin/EDTA), and gently dispose the excessive Trypsin/EDTA solution **within 30 seconds** with aspiration.
- Leave the T25 flask with the cells at **RT** for 3-5 minute (the cells usually will detach from the surface within 1-2 minutes). You must monitor the cells under microscope and when most of cells become rounded up, hit the flask gently against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.
- Add 5ml Trypsin Neutralization Buffer (cAP-28) and spin the cells down with 800g for 5 minutes.
- Re-suspend the cell pellet with 15ml of full cell culture medium and the cell suspension is transferred directly into 3 pre-coated T25 flasks (5ml each, and the cells are sub-cultured at 1:3 ratios)
- Change medium every 2-3 days and cells usually become confluent within 5-7days (when split at a 1:3 ratio).

NOTE: The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

Related Products:

Quick Coating Solution	cAP-01	240ml	Angio-Proteomie
Podocytes Growth Medium	cAP-44	500ml	Angio-Proteomie
Podocytes Basal Medium	cAP-44B	500ml	Angio-Proteomie
HBSS w/o Ca ²⁺ , Mg ²⁺	cAP-11	100ml	Angio-Proteomie
Cell Freezing Solution (FBS)	cAP-22	50ml	Angio-Proteomie
Cell Freezing Solution (Non-FBS)	cAP-22B	50ml	Angio-Proteomie
Trypsin/EDTA Solution	cAP-23	100ml	Angio-Proteomie
Trypsin Neutralization Solution	cAP-28	100ml	Angio-Proteomie
ITS (100x)	cAP-26	10ml	Angio-Proteomie
L-Glutamine-MAXIMUM (100x)	cAP-27	100ml	Angio-Proteomie
Human Plasma Fibronectin Solution	cAP-42	1mg/ml	Angio-Proteomie

THESE PRODUCTS ARE FOR RESEARCH USE ONLY

Caution: Handling human and animal tissue derived products is potentially bio-hazardous. Although each cell strain is tested negative for HIV, HBV and HCV DNA, or pathogens, diagnostic tests are not necessarily 100% accurate; therefore proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working with these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.