



Conditionally Immortalized Human Adipose-Derived Mesenchymal Stem Cells

ORDER INFORMATION

Name of Products: Conditionally Immortalized Human Adipose-Derived Mesenchymal Stem Cells (**CI-HAdMSCs**)
Catalogue Number: **cAP-0061**
Product Format: Frozen Vials
Cell Number: > 5 x 10⁵/vial
Viability: ≥ 70% when thawed from cryopreservation.

General Information

HAdMSCs (cAP-0051) are isolated from lipoaspirate derived from white adipose tissue and demonstrated with spindle-shaped-, and fibroblast-like cells. CI- HAdMSCs (cAP-0061) are selected from Puromycin resistant HAdMSCs after infected with lentiviruses expressing SV40 LT antigen under the control of TetON system. CI-HGAdMSCs are maintained with proliferative capacity in the presence of Doxycycline^{NOTE1} in Mesenchymal Stem Cell Growth Medium (MSCGM, cAP-35).

Characterization of the cells


Positive for CD29, CD44, CD73, CD90, CD105, and CD166 (greater than 95% of the cell population expresses these markers by flow cytometry).

Negative for CD14, CD31, CD34, and CD45 (less than 2% of cell population expresses these markers by flow cytometry).

CI-HBMMSCs are negative for HIV-1, -2, HBV, HCV, and Bacteria, Yeast and Mycoplasma.

Product Use: **CI-HAdMSCs** are for Research Use Only^{NOTE3}.

Shipping: Cells in Frozen Vials with Dry Ice Package.



cAP-0061 Conditionally Immortalized Human Adipose-Derived Mesenchymal Stem Cells with Potent Mitogen.

Handling of Arriving Cells

When you receive the cells in a frozen vial, you can transfer the vial of cells into a -80°C freezer for short-period storage or a liquid nitrogen tank for long-term storage. Thaw the cells in a 37°C water bath, and then quickly transfer the cells into a T75 flask with 15 ml MSCGM in the presence of 1ug/ml of Doxycycline and incubated overnight in a 37°C^{NOTE2} CO₂ incubator and change the medium next day (15 ml complete MSCGM) and every other thereafter.

Subculture Protocol:

- Rinse the cells in T75 flask with 15ml HBSS (**Room Temperature, RT**) twice.
- Add 4ml of Trypsin/EDTA (**RT**) (cAP-23) into one T75 flask (make sure the whole surface of the T75 flask is covered with Trypsin/EDTA), and gently dispose the excessive Trypsin/EDTA solution **within 30 seconds** with aspiration.
- Leave the T75 flask with the cells at **RT** for 1 minute (the cells usually will detach from the surface within 1-2 minutes). You can monitor the cells under microscope and when most of cells become rounded up, hit the flask against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.
- Add 10ml Trypsin Neutralization Buffer and spin the cells down with 800g for 5 minutes.
- Re-suspend the cell pellet with 30-45 ml of MSCGM in the presence of 1ug/ml Doxycycline and the cell suspension is transferred directly into 2 or 4 pre-coated T75 flasks (15ml each, and the cells are sub-cultured at 1:2 or 1:3 ratios)
- Change medium every 2-3 days and cells usually become confluent within 7 days (when split at a 1:3 ratio).

NOTE 1: To minimize the effect of SV40 antigen LT for your studies, cells should be cultured in the absence Doxycycline for 3-5 days.

NOTE 2: Although wild type SV40 LT antigen is used for cell immortalization, we noticed that the immortalized cells are growing better at 33-34°C, we encourage the end users to switch to 33-34°C if cells are growing slower at 37°C.

NOTE 3: The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

Related products

Quick Coating Solution	CAP-01	240ml	Angio-Proteomie
MSC Grown Medium (MSCGM)	CAP-35	500ml	Angio-Proteomie
MSC Basal Medium (MSCBM)	CAP-36	500ml	Angio-Proteomie
HBSS w/o Ca ²⁺ , Mg ²⁺	CAP-11	100ml	Angio-Proteomie
Trypsin/EDTA Solution	CAP-23	100ml	Angio-Proteomie
Trypsin Neutralization Solution	CAP-28	100ml	Angio-Proteomie

Human Material Precaution: All tissues used for isolation are obtained under informed consent and conform to HIPAA standards to protect the privacy of the donor's personal health information. It is best to use caution when handling any human cells. We recommend that all human cells be accorded the same level of biosafety consideration as cells known to carry HIV. With infectious virus assays or viral antigen assays, even a negative test result may leave open the possible existence of a latent viral genome.

Biosafety Level: 2

Angio-Proteomie Warranty: The viability of Angio-Proteomie' cell products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. Angio-Proteomie lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the Angio-Proteomie recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the Angio-Proteomie warranty for viability is no longer valid.

Caution: Handling human tissue derived products is potentially bio-hazardous. Although each cell strain is tested negative for HIV, HBV and HCV DNA, diagnostic tests are not necessarily 100% accurate; therefore, proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.