

## Goat anti-E2F7 Antibody

<b>Item Number</b>	dAP-1773
<b>Target Molecule</b>	Principle Name: E2F7; Official Symbol: E2F7; All Names and Symbols: E2F7; E2F transcription factor 7; FLJ12981; Accession Number (s): NP_976328.2; Human Gene ID(s): 144455; Non-Human GeneID(s):
<b>Immunogen</b>	DLSKQKKFTPERNP, is from internal region
<b>Applications</b>	Pep ELISA, WB Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 32000.
<b>Western Blot</b>	Western Blot: Approx 90kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 99.9kDa according to NP_976328.2). Recommended concentration: 0.3-1µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Li J, Ran C, Li E, Gordon F, Comstock G, Siddiqui H, Cleghorn W, Chen HZ, Kornacker K, Liu CG, Pandit SK, Khanizadeh M, Weinstein M, Leone G, de Bruin A. Synergistic function of E2F7 and E2F8 is essential for cell survival and embryonic development. Dev. Cell 2008 Jan 14 (1): 62-75..PMID:

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**